



Classified for
 "RESTRICTED USE"
 in New York State
 under 6NYCRR Part 326

Malice[®]
 75 WSP

ACCEPTED
 FOR REGISTRATION

Jan 7, 2010

New York State Department
 of Environmental Conservation
 Division of Solid & Hazardous Materials
 Pesticide Product Registration

Systemic and foliar insect control in turfgrass (including sod farms), and on fruit and nut trees, landscape
 ornamentals, and interior plantscapes and for control of listed insects infesting various crops.

ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl) methyl]-*N*-nitro-2-imidazolidinimine 75.0%

OTHER INGREDIENTS: 25.0%

TOTAL **100.0%**

Malice[®] 75 WSP contains imidacloprid, the active ingredient used in Merit[®] and Provado[®].

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

**EPA REG. NO. 34704-1009
 EPA EST. NO. 67545-AZ-001
 NET CONTENTS 4 x 1.6 OZ. WSP**

IHT 081109 V2D 12RB09

FORMULATED FOR


Loveland
 PRODUCTS INC.

P.O. BOX 1286
 GREELEY, COLORADO 80632-1286

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not do anything by mouth to an unconscious person
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

WPS USES: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses e.g., crops, sod farms, must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Protective eyewear
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

NON-WPS USES: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR part 170) – in general, only agricultural plant uses are covered by the WPS, must wear:

- Shirt and pants
- Gloves
- Protective eyewear
- Shoes plus socks

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Protective eyewear
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

- Keep children and pets off treated areas until dry.

RUNOFF MANAGEMENT

Do not cultivate within 10 feet of the aquatic areas to allow growth of vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

ENDANGERED SPECIES NOTICE

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

This product contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by this product and to other Group 4A products.

The active ingredient in this product is a member of the neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of this product and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Loveland Products, Inc. (LPI) strongly encourages the rotation to a block of applications with effective products of a different mode before using additional applications of neonicotinoid products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Foliar applications of this product or other Group 4A products from the neonicotinoid chemical class should not be used on crops previously treated with a long-residual, soil-applied products from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara®, Assail®, Calypso®, Centric®, Intruder®, Leverage®, Provado®, and Trimax™. Other 4A Group neonicotinoid products used as soil treatment include: Admire® and Platinum®.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

Restrictions:

- Keep children and pets off treated areas until dry.
- Do NOT apply through any type of irrigation system.
- Do NOT apply by air except for uses permitting aerial application in the "TREE, BRUSH AND VINE CROPS" section.
- Do NOT graze treated areas or use clippings from treated areas for feed or forage.
- Do NOT apply this product to soils that are waterlogged or saturated and avoid runoff or puddling of irrigation water following application.
- Do NOT allow leechate to run out for the first 10 days after application or reduced efficacy may result.
- Do NOT exceed a total of 8.6 oz./A per year (0.4 lb. a.i./A).

CROP USE LABEL PRODUCT PACKAGED IN WSP

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES AND COMMERCIAL FISH FARM PONDS.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Mixing and Loading Requirements

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

For Aerial Applications

Mount spray boom on the aircraft so as to minimize the drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Release spray at the lowest possible height consistent with good pest control and flight safety. Do not apply more than 10 feet above the crop canopy.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Because the potential for spray drift is high during temperature inversions, do NOT make aerial or ground applications during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Airblast (Air Assist) Specific Instructions for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e. turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

No-Spray Zone Requirements for Foliar Applications

Do not apply by ground within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

RESTRICTIONS

- DO NOT apply this product through any type of irrigation system.

- Unless specified within a crop specific application section for a given crop DO NOT apply more than 0.5 lbs. active ingredient per acre, per crop season regardless of formulation or method of application.

APPLICATION INSTRUCTIONS

Apply as a directed or broadcast foliar spray using adequate spray volumes, properly calibrated application equipment and spray adjuvant (if necessary) to obtain thorough coverage. Thorough coverage of the foliage (without runoff) is necessary. Loss of insect control or delay in onset of activity may result if there is not adequate coverage and retention of this product on leaves and fruit. Except where otherwise specified, this product may be applied using properly calibrated ground and/or aerial application equipment using a minimum recommended spray volume of 10 gallons/acre by ground application and 5 gallons/acre through aerial equipment.

Unless allowed under state-specific supplemental labeling, this product is not recommended for use on crops grown for production of true seed intended for private or commercial planting. Care must be taken to minimize exposure of this product to honey bees and other pollinators and use on crops requiring bee pollination must be avoided a minimum of 10 days prior to bloom through bloom. Your Cooperative Extension Service, PCAs, consultants or local LPI representatives can provide additional information on this product's uses with these crops.

Mixing Instructions

- Add a portion of the required amount of water to the spray tank and begin agitation.
- Add the recommended amount of this product.
- Fill the tank with the remaining water needed, being sure to maintain sufficient agitation during both mixing and application.
- If tank mixing this product with other pesticides and/or fertilizer solutions, please refer to the Compatibility Note below. When tank mixing this product with other pesticides, prepare the tank mixture as recommended above and follow the suggested Mixing Order below.

Mixing Order

Add this product first and allow the PVA packets to dissolve. Add any other wettable powders or wettable granules, flowables (suspension concentrates) second, and emulsifiable concentrates last. Maintain agitation as each component is added and do not add an additional component until the previous one is thoroughly mixed. A fertilizer pesticide compatibility agent may be needed if a fertilizer solution is added to the mix. To ensure uniformity of the spray mixture, be sure to maintain constant agitation during both mixing and application.

Compatibility Note

Do not use PVA packets in a tank-mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic that is not soluble in water or solvents. For further information, contact your local LPI representative.

Conduct the following test for compatibility of the intended tank mixture before adding this product to the spray or mix tank:

1. In a pint or quart jar, add proportionate amounts of each ingredient in the appropriate order.

2. Cap and shake for 5 minutes.
3. Let set for 5 minutes.
4. Observe the jar for signs indicating an incompatible mixture that should not be used such as poor mixing or the formation of precipitates that do not readily re-disperse.

Rotational Crops

As soon as practical following the last application, treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed.

Immediate Plant-back:

All crops on this label plus the following crops not on this label: barley, canola, corn (field, sweet and pop), rapeseed, sorghum, sugar beet and wheat.

30-Day Plant-back:

Cereals (including buckwheat, millet, oats, rice, rye and triticale), soybeans and safflower

10-Month Plant-back:

Onion and bulb vegetables

12- Month Plant-back:

All other crops

FIELD CROPS

Application Instructions

Apply as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage using the rates below is necessary to achieve optimum control. To improve coverage, a spray adjuvant may be used. This product may not knockdown heavy or established insect infestations; scout fields and make two applications if necessary to achieve control. For knockdown of pests or for improved control of other pests, this product may be tank mixed with other insecticides as recommended below.

COTTON

Do NOT graze treated fields after any application of this product

- Pre-Harvest Interval (PHI): 14 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per season: 6.5 ounces/A (0.31 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)
Banded-winged whitefly	0.7 – 1.3
Bollworm/Budworm (ovicidal effect)	
Cotton aphid	
Cotton fleahopper	
Green stink bug	
Plant bugs (excludes <i>Lygus hesparus</i>)	
Southern Green stink bug	
Lygus bug (<i>Lygus Hesperus</i>)†	1.0 – 1.3
Whiteflies (other than banded winged whitefly) †	

† Suppression only.

Tank Mix Recommendations

For Early-Season Control of Thrips:

- Mix 0.7 – 1.0 oz./A of this product with 1.6 – 3.2 oz./A of Bidrin® 8.

For Mid- to Late-Season Control of Cotton Leafperforator, Grasshoppers, Plant Bugs, Saltmarsh Caterpillar and Stink Bugs (Including Brown Stink Bug):

- Mix 0.7 – 1.0 oz. of this product with 4.0 – 8.0 oz. of Bidrin® 8 per acre.

Be sure to refer to the Bidrin® 8 label for specific use recommendations and to observe the most conservative use directions and precautions from both labels.

POTATO

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 4.0 oz./A (0.19 lb. a.i./A)

Pests Controlled**Application Rate (Oz./A)**

Aphids	1.0
Colorado potato beetle	
Flea beetles	
Leafhoppers	
Psyllids	

TOBACCO

- Pre-Harvest Interval (PHI): 14 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 6.0 oz./A (0.28 lb. a.i./A)

Pests Controlled**Application Rate (Oz./A)**

Aphids	0.5 – 1.1
Flea beetles	1.1
Japanese beetle	

VEGETABLE and SMALL FRUIT CROPS

Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website (www.epa.gov) for latest crop groups.

Application Instructions

Apply as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage using the rates below is necessary to achieve insect control. To improve coverage, a spray adjuvant may be used. This product may not knockdown heavy or established insect infestations; scout fields and make two applications if necessary to achieve control. For knockdown of pests or for improved control of other pests, this product may be tank mixed with other insecticides as recommended below.

FRUITING VEGETABLES

Crop Group 8 plus Okra including: Eggplant, Ground Cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet), Tomato, Pepinos, Tomatillo

NOT for use on crops grown for seed unless allowed by state-specific supplemental labeling.

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 5 days
- Maximum of this product allowed per crop season: 5.0 oz./A (0.23 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)	Application Instructions
Aphids Colorado potato beetle Leafhoppers Whiteflies	1.0	For insect control, good coverage of foliage and fruit is necessary. Incorporate applications of this product into a full-season program that uses effective products from multiple classes of chemistry and different modes of action in a blocked or windowed approach. For additional information, please contact your LPI representative, extension specialist or crop advisor.
Pepper weevil (Pepper only)	1.6	Apply the rate of this product using ground equipment only. Time applications prior to a damaging population becoming established.

GLOBE ARTICHOKE

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 14 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)
Aphids Leafhoppers	1.1 – 2.7

HEAD and STEM BRASSICA VEGETABLES

Crop Group 5 including: Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rage greens, Turnip (tops or leaves)

LEAFY VEGETABLES

Crop Group 4 including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (rocket), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chickory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only – Applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

NOT for use in California unless otherwise directed by supplemental labeling.

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum of this product allowed per crop season: 5.0 oz./A (0.23 lb. a.i./A)
- NOT for use on crops grown for seed unless allowed by state-specific supplemental labeling.

Pests Controlled	Application Rate (Oz./ A)	Application Instructions
Aphids Flea beetles Leafhoppers Whiteflies	1.0	For applications made to watercress: Apply to fully leafed-up canopies only. Production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application.

LEGUME VEGETABLES

Crops of Crop Group 6 (except soybean, dry) including:

Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

Bean (*Lupinus* spp. includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

Bean (*Phaseolus* spp. includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

Bean (*Vigna* spp. includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

Pea (*Pisum* spp. includes dwarf pea, edible pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

Other Beans and Peas: Broad bean (fava), chickpea (garbanzo bean), Guar, Jackbean, Lablab bean, hyacinth bean, lentil, pigeon pea, soybean (immature seed), sword bean

NOT for use on crops grown for seed unless allowed by state-specific supplemental labeling.

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 2.8 oz./A (0.13 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)
Aphids	0.9
Leafhoppers	
Whiteflies	

ROOT, TUBEROUS and CORM VEGETABLES

Crop Group 1 (except sugarbeet) including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden)[†], Burdock (edible)[†], Canna (edible, Queensland arrowroot), carrot[†], Cassava (bitter and sweet)[†], Celeriac[†], Chayote (root), Chervil (turnip-rooted)[†], Chickory[†], Chufa, Dasheen (taro)[†], Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip[†], Radish[†], Oriental radish (diakon)[†], Rutabaga[†], Salsify (black)[†], Salsify (oyster plant), Salsify (Spanish), Skirret, Sweet potato[†], Tanier (cocoyam)[†], Tumeric, Turnip[†], Yam bean (jicama, manioc pea), Yam (true)[†]

[†] Tops or green from these crops may be utilized for food of feed.

For recommended applications on Potato, refer to the Field Crops section.

NOT for use on crops grown for seed unless allowed by state-specific supplemental labeling.

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum of this product allowed per crop season: Radish: 0.9 oz./A (0.044 lb. a.i./A)
All other crops: 2.8 oz./A (0.13 lb. a.i./A)
- Maximum number of applications of this product per crop season: Radish: 1
All other crops: 3

Pests Controlled	Application Rate (Oz./A)
Aphids	0.9
Flea beetles	
Leafhoppers	
Whiteflies	

STRAWBERRY

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum of this product allowed per crop season: 3.0 oz./A (0.14 lb. a.i./A)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled**Application Rate (Oz./A)**

Aphids	1.0
Spittlebugs	
Whiteflies	

TREE, BUSH and VINE CROPS

Crops contained within certain crop groups recognized by EPA are subject to change. Refer to the EPA website (www.epa.gov) for latest crop groups

Application Instructions

Apply as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage using the rates below is necessary to achieve optimum control. To improve coverage, a spray adjuvant may be used. This product may not knockdown heavy or established insect infestations; scout fields and make two applications if necessary to achieve control. For knockdown of pests or for improved control of other pests, this product may be tank mixed with other insecticides as listed below.

Applying this product aerially may result in slower activity and reduced control relative to results from ground application.

For tree and vine crops, recommended application rates are based on full-size, mature trees or vines.

BUSH BERRY

Crop Subgroup 13 including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

- Pre-Harvest Interval (PHI): 3 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)
- Maximum number of applications of this product per crop season: 5
- Maximum application volume (water): 20.0 GPA – ground; 5.0 GPA – aerial

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled	Application Rate (Oz./A)
Aphids	0.8 – 1.1
Leafhoppers/Sharpshooters	
Blueberry maggot	1.6 – 2.1
Japanese beetles (adults)	
Thrips	

CITRUS

Crop Group 10 including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (Casimiroa spp.), and other cultivars and/or hybrids of these crops

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled	Application Rate (Oz./A)	Application Instructions
Aphids Asian citrus psyllid Black fly Leafhoppers / Sharpshooters Leafminers Mealybugs Scales Whiteflies	2.7 – 5.3 (depending on tree size, target pest and infestation pressure)	Scales: Time applications to the crawler stage and treat each generation.
Thrips†	2.7 – 5.3	

† Suppression only.

GRAPE

Including American bunch grape, Muscadine grape and Vinifera grape

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 14 days
- Maximum of this product allowed per crop season: 2.0 oz./A (0.1 lb. a.i./A)

Pests Controlled	Application Rate (Oz./ A)	Application Instructions
Leafhoppers / Sharpshooters Mealybugs	0.8 – 1.0	
Grapeleaf skeletonizer	1.0	Ground applications that provide thorough coverage of foliage should control grapeleaf skeletonizer. Aerial applications may provide suppression.

HOP

- Pre-Harvest Interval (PHI): 28 days
- Minimum interval between applications: 21 days
- Maximum of this product allowed per crop season: 6.4 oz./A (0.3 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)
Aphids	2.1

PECAN

NOT for use in California unless otherwise directed by supplemental labeling.

- Do NOT apply after shuck split.
- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 7.5 oz./A (0.35 lb. a.i./A)

Pests Controlled	Application Rate (Oz./A)
Aphids (use higher rate for Black pecan aphid) <i>Phylloxera</i> Spittlebugs	0.9 – 1.9

POME FRUIT

Crop Group 11 including: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled	Application Rate (Oz./ A)	Application Instructions
Leafhoppers	1.2 – 2.1	Apply low rate for low to moderate populations of white apple leafhoppers and high rate for high populations or for other leafhopper species. Apply this product while most leafhoppers are in the nymphal stage.
Aphids (except woolly apple aphid) Leafminers San Jose scale	2.1	Leafminer – To control first generation leafminer, apply as soon as pollination is complete and bees are removed from the orchard. Greatest control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. This product will not control late instar larvae. Rosy apple aphid – apply prior to leafrolling caused by rosy apple aphid. San Jose scale – time applications to the crawler stage. Treat each generation.
PEAR ONLY: Mealybugs Pear psylla	5.3	Mealybugs – apply maximum gallonage for tree with ground equipment. Ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of mealybugs.

STONE FRUIT

Crop Group 12 including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Apricot, Nectarine, Peach:

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 6.4 oz./A (0.3 lb. a.i./A)
- Minimum application volume (water): 50 GPA – ground application; 25 GPA – aerial application

Cherries, Plums, Plumcot, Prunes:

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)
- Minimum application volume (water): 50 GPA – ground application; 25 GPA – aerial application

Pests Controlled**Application Rate (Oz./A)**

Aphids	1.1 – 2.1
Green June beetle	
Japanese beetle	
Leafhoppers/Sharpshooters	
Plant bugs	
Rose chafer	
San Jose scale	
Cherry fruit fly (maggot of Eastern and Western)	1.6 – 2.1
Plum curculio†	
Stink bugs†	2.1

† Suppression only.

TROPICAL FRUIT

Including: Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Pulasan, Rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled	Application Rate (Oz./A)
Aphids	2.1
Leafhoppers/Sharpshooters	
Thrips	
Whiteflies	
Scales†	2.1

† Suppression only.

OTHER CROPS

Application Instructions

Apply as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage using the rates below is necessary to achieve optimum control. To improve coverage, a spray adjuvant may be used. This product may not knockdown heavy or established insect infestations; scout fields and make two applications if necessary to achieve control. For knockdown of pests or for improved control of other pests, this product may be tank mixed with other insecticides as listed below.

POPLAR/COTTONWOOD

Includes members of the genus *Populus* grown for pulp or timber

NOT for use in California unless otherwise directed by supplemental labeling.

- Minimum interval between applications: 10 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Pests Controlled	Application Rate (Oz./A)
Aphids	1.1 – 2.1
Leaf beetles	

CHRISTMAS TREE

- Minimum interval between applications: 7 days
- Maximum of this product allowed per crop season: 10.7 oz./A (0.5 lb. a.i./A)

Pests Controlled	Application Rate (Oz. / A)	Application Instructions
Aphids Adelgids Sawflies	1.1 – 2.1	Gall-forming adelgids – time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form, spraying will be ineffective.

TURF AND ORNAMENTALS FOR PRODUCT PACKAGED IN WSP

MIXING AND APPLICATION INSTRUCTIONS

Inside each foil pouch is a clear, water-soluble inner packet containing this product. To prepare a solution, remove the outer foil pouch and drop the required number of unopened clear water-soluble packets into the spray tank while filling with water to the desired level. Be sure to agitate while mixing and depending on the amount of agitation and the water temperature, the packets should completely dissolve within a few minutes of being added to the water. Note that cooler water temperatures increase the time needed for the inner packet to completely dissolve.

Mixing Precautions:

- Do NOT allow packets to become wet prior to adding to the tank.
- Do NOT handle the clean inner packets with wet hands or wet gloves.
- Do NOT use this product in a tank-mix with products that contain Boron or release free chlorine. Combining these products will result in a plastic that is not soluble in water or solvents (such as diesel oils, kerosene, gasoline or alcohol). Chlorinated water may be used.
- Because the water-soluble packets are not soluble in petroleum-based liquids, do NOT attempt to use this product's water-soluble packets directly in diesel oils or summer spray type oils such as those used in ULV or LV applications.
- Rough handling of the packets may cause breakage.
- Reseal outer carton to protect remaining packets.

Tank Mixes: This product has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. If this product is not known to be compatible with your particular tank mix partners, compatibility should be checked using the correct proportion of products in the following small jar test:

- 1) Add proportionate amount of each ingredient in the appropriate order to a pint or a quart jar;
- 2) Cap and shake for 5 minutes;
- 3) Let set for 5 minutes.

Poor mixing or formation of precipitates that do not readily re-disperse indicates an incompatible mixture that should not be used. For further information, contact your local LPI representative.

Mixing Instructions: The enclosed packets containing this product are water-soluble and will completely dissolve in water. The proper mixing procedure for this product alone or in tank mix combinations with other pesticides is:

1. Fill the spray tank $\frac{1}{4}$ to $\frac{1}{3}$ full with clean water.
2. While recirculating and with the agitator running, add the required number of unopened packets of this product.
3. The packets should completely dissolve in 5 to 10 minutes; allow sufficient time for thorough mixing.
4. Continue to fill spray tank with water until $\frac{1}{2}$ full.
5. If applicable, add remaining tank mix components in the following order: wettable powders, flowables, and emulsifiable concentrates. Ensure good agitation as each component is added. Do not add a tank mix component until the previous component is thoroughly mixed.
6. Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

TURFGRASS

This product will control listed soil-inhabiting pests in grassy areas such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made preceding or during the egg laying period. For insect control, make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Use Restrictions:

- Applications must NOT exceed a total of 8.6 oz. (0.4 lb of active ingredient) per acre per year.
- Applications must NOT be made when grassy areas are waterlogged or the soil is saturated with water because adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- Do not mow treated areas until after sufficient rainfall or irrigation has occurred in order to maintain the uniformity of the application.

Application Instructions:

Apply this product in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

Pest	Application Rate	Specific Instructions
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass atanius Cutworms† European Chafer European Crane Fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.	Grubs, European Crane Fly, billbugs and annual bluegrass weevil: For best results make applications prior to egg hatch of the target pest. Chinchbugs: Make applications prior to the hatching of the first instar nymphs. Mole Crickets: Make applications prior to or during the peak egg hatching period. When adults or large nymphs are present and actively tunneling, this product should be accompanied by a curative insecticide. For insect control, the active ingredient must be moved through the thatch by irrigation or rainfall occurring within 24 hours after application.
Chinchbugs† Mole Crickets	1.6 oz. (1 packet) per 8,250 sq. ft.	

ORNAMENTALS, GROUNDCOVERS AND INTERIOR PLANTSCAPES

This product is a systemic insecticide that may be applied to ornamentals, groundcovers and interior plantscapes in and around industrial and commercial buildings and residential areas. The insecticide is translocated upward into the plant system and for best results must be placed where the growing portions of the target plant can absorb the active ingredient. When applicable, adding a fertilizer containing nitrogen into the spray solution may enhance plant uptake of this product.

Rotational Crops:

As soon as practical following the last application, treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed. Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed.

Ant Management Programs:

This product may be used to limit the honeydew available as a food source for ant populations when controlling aphids, scale insects, mealy bugs and other sucking pests on ornamentals. Applications of this product may be supplemented with bait traps, residual sprays and other methods to further reduce the unwanted ant population.

Insect Resistance:

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

Woody Perennials:

Protection in woody perennials is slower than in herbaceous species and a delay of 2 or more weeks should be expected, with longer delays for larger plants. Because of this, applications to woody perennials should be made well in advance of expected insect activity.

Bark Media:

Treatments of this product to media with 30 - 50% or more bark content may confer a shorter period of protection.

FOLIAR AND BROADCAST APPLICATIONS

This product may be applied as a broadcast or foliar application to trees (including non-bearing fruit and nut trees), shrubs, evergreens, flowers, foliage plants, ground covers, interior plantscapes and vegetable plants intended for resale.

Application Instructions:

Apply this product in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

When making foliar applications to plants with hard-to-wet foliage such as holly, pine or ivy, use of a spreader/sticker is recommended.

Pest	Application Method	Application Rate	Specific Instructions
Adelgids Aphids Japanese beetle (adult) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy- winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips† Whiteflies	Foliar	1.6 oz. (1 packet) per 300 gal. of water	Make applications prior to establishment of large pest populations and retreat as necessary. Applying this product foliarly after a soil application in the same crop is not recommended for resistance management purposes.
White grub larvae (such as Japanese beetle larvae, chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle and Oriental beetle)	Broadcast	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.	Mix the specified amount of this product in sufficient water to uniformly cover the area being treated using at least 2 gallons of water per 1000 sq. ft. For insect control, incorporate this product into the upper soil profile by irrigating after the application is made.

† Suppression only.

SOIL INJECTION AND DRENCH APPLICATIONS

Application Site	Application Rate	Application Instructions	Pests Controlled
<p>Trees (Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.)</p>	<p>1.6 oz. (1 packet) per 24 – 48 in. of cumulative trunk diameter (DBH)</p>	<p>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York. GRID SYSTEM: Holes must be spaced on 2.5 ft centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For insect control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p>SOIL DRENCH: Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.</p>	<p>Adelgids Aphids Armored Scale† Black vine weevil larvae Eucalyptus Longhorned Borers Emerald Ash Borer Flatheaded Borers (including bronze birch and alder borers) Japanese Beetles (adults) Lacebugs Leaf Beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips† White grub larvae Whiteflies</p>
<p>Shrubs</p>	<p>1.6 oz. (1 packet) per 24 – 48 ft. of cumulative shrub height</p>	<p>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Using a minimum of 4 holes per shrub, apply to individual plants maintaining a low pressure and use sufficient</p>	<p>Adelgids Aphids Armored Scale† Black vine weevil larvae Eucalyptus Longhorned Borers Emerald Ash Borer Flatheaded Borers (including bronze birch and alder borers) Japanese Beetles (adults) Lacebugs Leaf Beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips† White grub larvae Whiteflies</p>

Table cont'd. next page

SOIL INJECTION AND DRENCH APPLICATIONS

Application Site	Application Rate	Application Instructions	Pests Controlled
<i>Shrubs cont'd.</i>	1.6 oz. (1 packet) per 24 – 48 ft. of cumulative shrub height	<p>solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days.</p> <p>SOIL DRENCH: Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.</p>	
Flowers and Ground Cover	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, irrigate thoroughly after application.	

† Suppression only of these species

POME FRUIT IN AND ON RESIDENTIAL AREAS

Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear) Quince

- Pre-Harvest Interval (PHI): 7 days
- Reapplication Interval: At least 10 days
- Maximum applications per year: 5

Pest	Ounces per 300 Gals. of Water	Ounces per Acre‡	Specific Instructions
<p>Aphids (except Woolly apple aphid)</p> <p>Leafhoppers (including glassy-winged sharpshooter)</p> <p>Leafminer</p> <p>Mealybugs†</p> <p>San Jose Scale†</p>	1.6 (1 packet)	2.1	<p>Apply as a foliar spray as needed after petal-fall is complete.</p> <p>Rosy Apple Aphid: Apply prior to leaf rolling caused by the pest.</p> <p>Leafhopper: For late season (preharvest) control, apply while most leafhoppers are in the nymphal stage.</p> <p>Leafminer: Make first application as soon as petal-fall is complete for control of first generation, with best results occurring when the application is made at the earliest possible time.</p>

Table cont'd. next page

Pest	Ounces per 300 Gals. of Water	Ounces per Acre‡	Specific Instructions
<i>Aphids (except Woolly apple aphid)</i> <i>Leafhoppers (including glassy-winged sharpshooter)</i> <i>Leafminer</i> <i>Mealybug†</i> <i>San Jose Scale† cont'd.</i>	1.6 (1 packet)	2.1	For succeeding generations, best results occur when applications are made early in the adult flight against egg and early instar larvae. If generations are overlapping or severe pressure continues, a second application may be necessary after 10 days. A single application may result in suppression only. This product will not control late stage larvae. Mealybug: For insect control be sure to have good spray coverage of the trunk and scaffolding limbs or other nesting sites. San Jose Scale: Time applications to the crawler stage and treat each generation.

† Not permitted for control on pears in California.

‡ The amount of this product required per acre depends on tree size and volume of foliage. The listed rate per acre is based on a standard of 400 gallons of dilute spray per acre for large trees.

PECANS IN AND ON RESIDENTIAL AREAS

NOT PERMITTED IN CALIFORNIA unless otherwise directed by supplemental labeling

- Reapplication Interval: At least 10 days
- Maximum applications per year: 3
- Maximum of this product allowed per year: 6.3 oz./A

Pest	Ounces per 300 Gals. of Water	Ounces per Acre‡	Specific Instructions
Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1.6 (1 packet)	2.1	Apply as a foliar spray as pest pressure builds but before infestation is extremely heavy. Two applications at a 10 – 14 day interval may be required to achieve control. For insect control, thorough and uniform coverage is necessary. Coverage may be improved through the use of an organosilicone-based spray adjuvant.

‡ The amount of this product required per acre depends on tree size and volume of foliage. The listed rate per acre is based on a standard of 400 gallons of dilute spray per acre for large trees.

GRAPES ORNAMENTAL USE

- Reapplication Interval: At least 14 days
- Maximum of this product allowed per year: 2.0 oz./A

Ornamental Grapes In Industrial and Commercial Buildings and On Residential Areas

Pest	Ounces per 300 Gals. of Water	Ounces per Acre	Specific Instructions
Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.6 (1 packet)	1.0	Apply as a foliar spray using 200 gallons of water per acre.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE [For product packaged in plastic containers]: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE STORAGE [For product packaged in Water-soluble Packaging]: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Exposure to moisture or excessive handling of water-soluble packets may cause breakage.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse or refill this container.

Outer Packaging: Outer Packaging for this product is secondary packaging to contain either non water soluble plastic bags or water soluble plastic bags: Thoroughly rinse any soluble powder residue from pail or box into application equipment; then offer for recycling if available or dispose of in a sanitary landfill.

Non water-soluble plastic bags: Completely empty plastic bags into application equipment, ensure that all product is removed from the bag and then offer for recycling if available or dispose of in a sanitary landfill.

Water-soluble plastic bags: After adding water-soluble plastic bags to the spray tank, allow sufficient time for bags to dissolve before spraying. There is no container disposal once the bag has been dissolved in the spray tank.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE. IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Actara®, Centric®, and Platinum® are registered trademarks of a Syngenta Group Company.

Admire®, Calypso®, Leverage®, Merit®, and Provado® are registered trademarks and Trimax™ is a trademark of Bayer.

Assail® is a registered trademark of Nippon Soda Co., Ltd. and Intruder® is a registered trademark of Nippon Soda Co., Ltd. and licensed for distribution by E.I. duPont de Nemours & Company.

Biden® is a registered trademark of AMVAC Chemical Corporation.

Malice® is a registered trademark of Loveland Products, Inc.

Bidrin and Leverage are restricted use pesticides.